



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Elementary Proof of Cauchy's Theorem.

BY ARTHUR LATHAM BAKER.

Denote the sides of a right triangle (Argand Diagram) by x, iy, z . Multiply each of these by the complex number w (plane vector) giving $wz = wx + wiy$. Hence the proposition: *If on the three sides of a right triangle, similar and similarly placed triangles be constructed, then the sum of the corresponding sides (considered as vectors) of the leg triangles is equal to the corresponding side of the hypotenuse triangle.*

At the limit this becomes

$$dW = wdz = wdx + widy = w(dx + idy).$$

Hence *the change in the function W is the same whether z follows the elementary paths dx, idy , or the resultant of these, dz .*

Hence $W = \int dW$, the sum of all these changes in W due to changes in z , will be the same whether z follows the elemental paths $\Sigma dx, \Sigma idy$ or resultants of these, so long as in the deformation of the path of z into Σdx and Σidy we do not pass over any point in which $w = \infty$.

Hence W *always attains the same value for the same value of z , whatever the path of z , provided no point where $w = \infty$ is enclosed between the paths.*

Since we can take the end of the path of z as near the beginning as we choose, or coincident with it, we can say:

(Cauchy's Theorem) $W = \int dW = \int wdz$ *taken around a closed curve enclosing no point where $w = \infty$ is zero.*

UNIVERSITY OF ROCHESTER.